BioRisk

BioRisk 24: 55–67 (2026)
DOI: 10.3897/biorisk.24.180507

Short Communication

# Green guardians: Mosses as potential low-cost biomonitors of air pollution in urban Chicago

Thomas Campbell[1], Gilbert F. Gwilliam III[2], Nkosi Michael Evans[1,3], Kate Peltz[1], Glenn Simpson[1], Laure Dussubieux[4], Emily Erazo[1], Kage Guare[1], Agnetta Krechner[1], Estefania Long[1], Stephanie Maari[2], Isaac Oguike[1], Kimberly Rice[1], Amanda Zinke[1], Chrissy Christian[2], Ayesha Qazi-Lampert[2], Joshua M. Henkin[5,6], Danny Kreider[2,7], Christopher Maves[2], Gary Merrill[2], Phoebe Moy[2,8], Yarency Rodriguez[2], Zoe Ryan[3,7], Matt von Konrat[2]

1 *Biology Department, College of Arts and Sciences, Northeastern Illinois University, Chicago, Illinois, USA*
2 *Gantz Family Collections Center, Science & Education, Field Museum of Natural History, Chicago, Illinois, USA*
3 *Department of Botany, University of Wisconsin, Madison, Wisconsin, USA*
4 *Negaunee Integrative Research Center, Science & Education, Field Museum of Natural History, Chicago, Illinois, USA*
5 *Analytics Lounge, 1052 N Milwaukee Ave FL2, Chicago, Illinois, USA*
6 *Department of Plant Biology, Rutgers University, New Brunswick, New Jersey, USA*
7 *University of Illinois, Chicago, Illinois, USA*
8 *DePaul University, Chicago, Illinois, USA*
*Corresponding author: Matt von Konrat (mkonrat@fieldmuseum.org)*

OPEN ACCESS

Academic editor: Josef Settele
Received: 26 November 2025
Accepted: 19 January 2026
Published: 30 March 2026

Citation: Campbell T, Gwilliam III GF, Evans NM, Peltz K, Simpson G, Dussubieux L, Erazo E, Guare K, Krechner A, Long E, Maari S, Oguike I, Rice K, Zinke A, Christian C, Qazi-Lampert A, Henkin JM, Kreider D, Maves C, Merrill G, Moy P, Rodriguez Y, Ryan Z, von Konrat M (2026) Green guardians: Mosses as potential low-cost biomonitors of air pollution in urban Chicago. BioRisk 24: 55–67. https://doi.org/10.3897/biorisk.24.180507

Copyright: © Thomas Campbell et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

## Abstract

According to the Lancet Commission, air pollution is one of the single largest environmental causes of premature death in the world. Particulate matter (PM), including metals, is among the pollutants of concern. Bryophytes, especially mosses, have been used for many decades as low-cost biomonitors of environmental deposition of heavy metals. In this exploratory investigation, we use strategically placed *Sphagnum* moss paired with portable X-ray fluorescence (pXRF) analysis as a living biomonitor for air/particulate metal pollution in and around the city of Chicago, Illinois, U.S.A. Preliminary results reveal a correlation between urbanization and some metal particulate matter, especially iron. The study warrants further investigation and supports the use of mosses and pXRF spectrometry as a rapid, low per-sample cost, and potentially accurate alternative to conventional metal pollution monitoring.

**Key words:** Air pollution, biological monitoring, bryophytes, Chicago, heavy metals, mosses, particulate matter, pXRF, *Sphagnum*

## Introduction

The impact of air pollution represents one of the most pressing environmental health challenges of our time and has been well documented (Chen and Kan 2008; Sharma et al. 2013; Manisalidis et al. 2020; Fuller et al. 2022; Shleag et al. 2024). The 2018 Lancet Commission on pollution and health identified air pollution as the single largest environmental cause of disease and premature death globally, accounting for 16% of all deaths (Landrigan et al. 2018). Recent studies have established connections between air pollution expo-

sure and COVID-19 severity (Liang et al. 2020; Travaglio et al. 2021). Urban environments are particularly vulnerable to air pollution, with cities serving as potential hotspots for air pollution-related diseases (Nieuwenhuijsen 2020). The exact extent of health effects at the city level from urban air pollution remains largely unknown (Khomenko et al. 2021). This lack of knowledge is particularly alarming in the United States, given that it is among the most urbanized countries globally; in 2011 alone, urbanization had sprawled 11% relative to 2001 (Bounoua et al. 2018).

Air pollution comprises a complex mixture of gaseous and particulate components, each potentially harmful to the cardiovascular and respiratory systems (Hamanaka and Mutlu 2018). Particulate matter (PM) represents a critical component in air pollution with well-documented health impacts (Aryal et al. 2021). Common components of PM include nitrates, sulfates, polycyclic aromatic hydrocarbons, endotoxins, and heavy metals such as iron (Fe), copper (Cu), nickel (Ni), zinc (Zn), and vanadium (V) (Brook et al. 2010; Newby et al. 2015). We note that the term "heavy metal" has been broadly applied in the plant sciences, but not without controversy, since the term "heavy metal" seems to suggest that these metals or their compounds are toxic (Appenroth 2010). Heavy metals (HMs) specifically are emitted into the environment from common urban sources such as transportation, industry, and urbanization, as well as from fossil fuels and agriculture (Koz et al. 2008). Specifically, transportation has been shown to contribute Cu and Zn due to wear of brake linings (Liu et al. 2023).

Biomonitoring is generally defined as the systematic use of living organisms or their responses to determine the condition or changes in the environment (Thomas 1961). Biological monitoring is an important tool for evaluating the negative impacts of human activities on the atmosphere (Alam 2018). Green land plants, including bryophytes, have been effectively used as air pollution indicators for many years (Badamasi 2017). Bryophytes are of great ecological and environmental significance, playing an important role as possible indicators of climate change (Lindo et al. 2013; Ruklani et al. 2021), in nutrient cycling (Rieley et al. 1979), and, through their water retention, in reducing soil nutrient loss and flooding risk (Anderson et al. 2010). Bryophytes also serve as the "macrophytes," providing a matrix in which many microscopic organisms live, including tardigrades, mites, rotifers, micro-mollusks, microalgae, microfungi, and prokaryotes (Gerson 1982; Huttunen et al. 2017).

Mosses, in particular, have been used for more than five decades to estimate atmospheric heavy metal deposition (Ruehling and Tyler 1968). Similarly, liverworts have been used as biomonitors of heavy metal pollution (Basile et al. 2013; Postiglione et al. 2025). Since 1990, an ongoing moss survey has had an important role in identifying spatial and temporal trends in atmospheric heavy metal pollution across Europe (Harmens et al. 2010). Bryophytes, and mosses in particular, have been shown to be effective low-cost biological monitors of metal air pollution because they readily accumulate particulate matter over time, reflecting long-term pollution levels (Fernández et al. 2015; Messager et al. 2021). Bryophytes have a number of morphological and physiological properties that make them ideal bioindicators. These have been extensively reviewed by Blagnytė and Paliulis (2010), for example, their relatively small size, their ability to survive in highly polluted areas, the lack of roots and ac-

quisition of nutrients needed for vital processes from wet and dry deposition, and resistance to many substances that are highly toxic to other organisms. Interestingly, mosses have also been used as potentially useful biomonitors for the atmospheric deposition of anthropogenic microfibers and microplastics (Roblin and Aherne 2020).

There are two well-established types of biomonitoring using mosses to evaluate atmospheric contamination: 1) passive biomonitoring, using moss that grows naturally in a particular area, and 2) active biomonitoring, by transplanting moss from other locations (Ares et al. 2012). Spectroscopic techniques used to determine the concentration of particulate matter in moss samples require expensive instrumentation and involve time-consuming sample preparation protocols with heavy use of reagents (Messager et al. 2021). In situ and laboratory X-ray fluorescence (XRF) spectrometry of mosses has been presented as a rapid, low-cost, and accurate alternative to conventional metal pollution biomonitoring (Koz et al. 2008; Šoltés and Gregušková 2013; Messager et al. 2021).

Here, we present an exploratory investigation into the potential application of mosses (*Sphagnum*) as a biological monitor for particulate matter in Chicago, Illinois, U.S.A., using pXRF spectrometry analysis. The aims of our study include 1) to provide an initial evaluation of the potential use of mosses as biological indicators for air pollution, particularly heavy metals; 2) to explore whether pXRF spectrometry is an effective screening mechanism that warrants further study; and 3) to provide a preliminary investigation into different laboratory protocols and material preparations.

## Materials and methods

### Early pilot studies testing feasibility

Between 2017 and 2024, we conducted a series of preliminary pilot studies to assess the feasibility of using portable X-ray fluorescence (pXRF) spectrometry and mosses as biomonitors of atmospheric heavy metal deposition. These early efforts included in situ sampling across Chicago, testing various handheld and benchtop pXRF units, testing various species of moss, and deploying *Sphagnum* moss bags in collaboration with local volunteers. While these initial investigations were exploratory and varied in methodology, they laid the foundation for the standardized, systematic approach described in this paper. Additional details and results from these earlier efforts are included as supplemental material to recognize contributors and provide context for the evolution of the project (see https://zenodo.org/records/15097659 and Data availability below).

### Moss preparation and processing

We used two primary sources of *Sphagnum*: 1) commercial *Sphagnum* moss (*Sphagnum squarrosum*), purchased from Moss Acres (Honesdale, PA, USA) in June 2024; and 2) *Sphagnum fallax* from Maine, collected by Blanka Shaw in June/July 2023 and 2024. Supplementary files, including experimental detail, data, and presentations, are available at https://zenodo.org/records/15097659. The voucher is available at the Field Museum and

can be accessed here (https://collections-botany.fieldmuseum.org/catalogue/4871426). The moss was carefully cleaned to remove invertebrates, debris, and non-moss species, retaining only bright green segments. Rhizomes and pale portions were manually removed and discarded. Samples were air-dried overnight on newspaper, followed by screening through a grated mesh to remove remaining particulate matter.

## Sample washing protocol

Three sequential washing treatments were applied using 500 mL of Chicago tap water in aluminum trays. Moss samples were soaked and gently agitated for 5 minutes in each tray, with excess water removed between treatments. Following the final wash, remaining debris was removed with forceps, and samples were air-dried for 48 hours.

## Moss bag preparation

Standardized 1 g samples of prepared *Sphagnum* were distributed into 7.5 × 10 cm mesh bags with drawstring closures. Each bag received a unique identifier and was attached to a 20 cm hanging string with a plastic plant label secured by a binder clip.

## Site selection and deployment

Distribution sites were selected based on accessibility and safety considerations, representing hypothesized potential pollution sources, including 1) an industrial point source (a metals recycling facility); 2) an industrial/transportation corridor (Indian Ridge Marsh); 3) a transportation hub (Des Plaines railroad station); 4) a semi-natural reference area (a Lake County natural area) (Fig. 1A); and 5) a vehicle exhaust study (moss bags attached to a 2006 Ford F-350 diesel truck) (Fig. 1B). Control samples were maintained at the Field Museum under both sealed (plastic bag) and exposed conditions. Deployment occurred during June–July 2024, with collection after 1-, 3-, and 4-week exposure periods.

## pXRF analysis

Retrieved moss samples were processed using liquid nitrogen in a clean mortar and pestle to create a homogeneous powder. Powdered samples were packed into labeled pXRF sample cups and sealed with a Prolene thin-film membrane to prevent contamination. Analysis was conducted using a Tracer 5 Bruker pXRF instrument with identical settings across all samples. Target elements included titanium (Ti), manganese (Mn), iron (Fe), nickel (Ni), copper (Cu), and zinc (Zn). Multiple measurements were performed on each sample to improve data reliability.

Reference samples, including those washed in different water sources and those left unwashed, were analyzed to compare the effects of the washing treatments on metal levels. Detailed settings and procedures used to operate the pXRF unit are extensively outlined at https://zenodo.org/records/15097659 and Data availability below.

**Figure 1. A.** Biobags attached to branches in a semi-natural reference area; **B.** Biobags taped to the exhaust of a Ford F-350 diesel truck.

### Data analysis

Results are reported as ratios of exposed sample counts to unexposed sample counts ("ratio of counts/control"). Six replicate analyses were performed on 2–3 moss bags from each location, providing 12–18 measurements per element per location. Given that there were some differences in metal content between the two sources of moss, we kept them separate and analyzed results of exposed moss relative to reference samples from the same moss sources.

## Results

### Atmospheric heavy metal capture

Twelve moss bags were deployed across four Chicagoland locations during June and July 2024. After 25–30 days of exposure, pXRF analysis revealed location-specific metal accumulation patterns compared to reference samples (Fig. 2). The metals recycling facility (site 1) showed elevated levels across all measured elements, consistent with its role as a multi-metal emission source. Indian Ridge Marsh (site 2), situated adjacent to industrial and transportation facilities, demonstrated significantly increased iron and nickel concentrations. The Des Plaines railroad station (site 3) exhibited elevated iron levels, potentially reflecting particulate generation from steel rail infrastructure. Notably, even the semi-natural site in Lake County (site 4) showed elevated nickel concentrations, likely attributable to its proximity (<1 km) to a major highway, potentially demonstrating the far-reaching influence of transportation-related emissions.

### Vehicle emission detection

Six moss bags attached to a 2006 diesel Ford F-350 truck operating in the Chicago area showed time-dependent metal accumulation patterns (Fig. 3). After three days of exposure, significant increases were observed in iron, zinc, copper, titanium, and manganese compared to controls. A reduction in metal levels was observed after four days of exposure, which coincided with significant rainfall in the Chicago area, suggesting potential washing effects on accumulated particulates. This observation highlights the importance of considering meteorological factors in biomonitoring protocols.




**Figure 2.** Ratio of pXRF counts for exposed samples to pXRF counts for unexposed samples for six metals. The horizontal line represents a ratio of 1, i.e., no net accumulation of that metal in the exposed samples. Error bars represent the standard deviation of the ratio of the measurements of the exposed samples to the unexposed samples.




**Figure 3.** Ratio of pXRF counts for exposed samples to pXRF counts for unexposed samples for six metals versus exposure time of the samples.

## Discussion

This exploratory study demonstrates the feasibility of using standardized *Sphagnum* moss bags coupled with pXRF analysis for monitoring atmospheric particulate matter in urban environments. Our results align with a growing body of evidence supporting this technique as a cost-effective tool for preliminary environmental contamination assessments, providing rapid, low per-sample cost, and accurate alternatives to conventional metal pollution biomonitoring (Fernández et al. 2015; Messager et al. 2021).

### Heavy metal sources and patterns

The metal accumulation patterns observed in our study are consistent with known urban emission sources. Elevated iron levels across multiple sites likely reflect its widespread presence in urban particulate matter, predominantly originating from vehicle exhaust, brake wear, rail transport, and industrial processes (Abbasi et al. 2013; Mohsen et al. 2018; Mayer et al. 2024). The high iron recorded near railroad infrastructure supports previous studies linking rail transport with significant metal particulate emissions (Abbasi et al. 2013; Mohsen et al. 2018). Copper and zinc accumulation in mosses exposed to vehicle emissions reinforces established associations of these metals with transportation sources, particularly from brake lining abrasion and tire wear (Pearson et al. 2000; Mayer et al. 2024). Elevated nickel concentrations detected at the semi-natural Lake County site suggest the possibility of regional atmospheric metal dispersion potentially linked to transportation-related emissions, a pattern observed in European moss monitoring studies (Godzik 2020; Vegetation ICP 2020). Interestingly, recent μ-SRXRF analyses provided insights into how heavy metals distribute at the microscale within moss tissues, enhancing understanding of accumulation dynamics observed in *Sphagnum* bags (Weinberger et al. 2025). Similarly, moss biomonitoring has been successfully applied to map detailed spatial patterns of metal contamination in urban environments, underscoring the utility of moss bags in pollution assessments (Urošević et al. 2023).

### Methodological considerations

pXRF spectrometry proved valuable for rapid, cost-effective screening of metal concentrations in moss samples. The ratio-based analytical approach effectively normalized variation between moss sources, providing meaningful comparisons across sites and exposure periods. While our washing protocol appeared adequate in removing surface contamination and maintaining moss bioaccumulation capacity, future standardization of sample preparation—particularly washing duration, water quality specifications, and drying protocols—would enhance reproducibility and comparability of results (Fernández et al. 2015; Messager et al. 2021). Temporal dynamics observed in vehicle-mounted samples emphasize the need for standardized exposure durations and incorporation of meteorological data. The observed decrease in accumulated metals after rainfall underscores the significant impact weather conditions have on biomonitoring outcomes and highlights the importance of developing moss bag designs that mitigate wash-off effects.

## Broader implications

Our study addresses a notable research gap in North American bryophyte biomonitoring, complementing the established European moss survey network (Vegetation ICP 2020). Demonstrating the effective application of moss biomonitoring in Chicago indicates strong potential for broader adoption across other US urban regions. Moss bag biomonitoring offers a cost-effective, community-accessible alternative to laboratory-based monitoring, enabling resource-limited communities disproportionately affected by pollution to document environmental contamination and advocate for remediation. Combining moss biomonitoring with portable XRF technology can facilitate community-based environmental monitoring, enabling frequent and widespread pollution assessments at significantly reduced costs (Messager et al. 2021).

## Study limitations

Several limitations must be recognized. The exploratory nature of this study involved limited sample sizes and temporal coverage. Although weather impacts were noted, systematic meteorological evaluations were not conducted. Additionally, commercially sourced *Sphagnum* moss, while facilitating standardization, may introduce baseline contamination variability compared to locally sourced material (Fernández et al. 2015). Other potential sources of contamination include the water used to wash the mosses and the newspaper on which they were dried. Furthermore, pXRF provides semi-quantitative results and may have detection limits unsuitable for low-level contamination scenarios (Šoltés and Gregušková 2013; Messager et al. 2021).

## Future research directions

This preliminary study suggests multiple avenues for future research. Replication of established moss biomonitoring protocols would enhance methodological consistency, allowing broader comparative analyses (Fernández et al. 2015; Vegetation ICP 2020). Expanding this approach to additional urban settings and pollution sources could elucidate broader regional pollution patterns and facilitate comprehensive air quality assessments.

Integrating meteorological data and examining seasonal variations would improve understanding of temporal dynamics in metal accumulation. Direct comparisons between active (transplanted) and passive (native) moss monitoring techniques could further optimize deployment strategies to achieve specific environmental monitoring objectives (Ares et al. 2012).

Utilizing historical herbarium moss specimens for retrospective pXRF analyses presents a compelling opportunity for reconstructing historical pollution profiles, evaluating long-term environmental changes, and assessing remediation effectiveness (Belloeil et al. 2021). Emphasizing community-based biomonitoring initiatives could significantly enhance environmental justice efforts by providing accessible tools for marginalized communities facing disproportionate pollution exposure (Fuller et al. 2022).

## Conclusion

This study highlights the practicality and effectiveness of using *Sphagnum* moss coupled with pXRF analysis for urban atmospheric pollution biomonitoring. This approach offers significant promise as a cost-effective and accessible environmental monitoring tool. Future research, including broader geographical application and methodological standardization, will strengthen its utility in addressing environmental justice and promoting sustainable urban air quality management.

## Acknowledgments

We would like to thank the Field Museum's Women's Board for financial support, as well as Dr. Lesley de Souza for coordinating the Women in Science Internship Program. Several students were financially supported by Northeastern Illinois University's Student Center for Science Engagement. We also extend our gratitude to Alizé Gonzalez for assistance with material and sample collection and processing. The generous support of the National Science Foundation (Award Nos. 1145898, 1146168, 2001509) is gratefully acknowledged. The iDigBio Natural History Collections Summer Internship Program and the DePaul University Dean's Undergraduate Fellowship are also acknowledged. The portable XRF used in this study is part of the Elemental Analysis Facility at the Field Museum and was acquired through the generosity of the Grainger Foundation.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Use of AI

During the conceptualization phase for the supplementary file's structure and appropriate potential structure for methodology based on lab notes, ChatGPT [OpenAI. (2025). ChatGPT (GPT-5.1) [Large language model]. https://www.openai.com/chatgpt] was used to brainstorm organizational ideas. The AI-generated suggestions served as a starting point and were subsequently reviewed, edited, and refined, and full responsibility is taken for the final structure and content. Prompts used included: "can you produce a two page summary","can you produce a one page summary of the attached for the methods section of a scientific report" based on a PDF of lab notes taken throughout 2024.

### Funding

We further recognize the support of the Grainger Bioinformatics Center, as well as curatorial assistance provided by L. Kawasaki, D. Le, A. Sreniawski, and A. Balla (Field Museum). We are grateful to Peter Zieba, Analytics Lounge, Chicago, Illinois, United States of America, for technical support during the early phases of this project.

## Author contributions

All authors have contributed equally.

## Author ORCIDs

Thomas Campbell https://orcid.org/0000-0003-4228-9685
Gilbert F. Gwilliam III https://orcid.org/0009-0006-4072-3539
Stephanie Maari https://orcid.org/0009-0008-6772-8203
Chrissy Christian https://orcid.org/0009-0009-1293-5931
Ayesha Qazi-Lampert https://orcid.org/0000-0002-5733-0911
Yarency Rodriguez https://orcid.org/0009-0009-4253-8355

## Data availability

All data and procedures used in the preparation of this paper are available at https://doi.org/10.5281/zenodo.15097659.

## References

Abbasi S, Jansson A, Sellgren U, Olofsson U (2013) Particle emissions from rail traffic: A literature review. Critical Reviews in Environmental Science and Technology 43: 2511–2544. https://doi.org/10.1080/10643389.2012.685348

Alam A (2018) Bryomonitoring of environmental pollution. In: Vats S (Ed.) Biotic and Abiotic Stress Tolerance in Plants. Springer Singapore, Singapore, 349–366. https://doi.org/10.1007/978-981-10-9029-5_13

Anderson M, Lambrinos J, Schroll E (2010) The potential value of mosses for stormwater management in urban environments. Urban Ecosystems 13: 319–332. https://doi.org/10.1007/s11252-010-0121-z

Appenroth K-J (2010) Definition of "Heavy Metals" and Their Role in Biological Systems. Soil Heavy Metals. Soil Biology. Springer Berlin Heidelberg, Berlin, Heidelberg, 19–29. https://doi.org/10.1007/978-3-642-02436-8_2

Ares A, Aboal JR, Carballeira A, Giordano S, Adamo P, Fernández JA (2012) Moss bag biomonitoring: A methodological review. The Science of the Total Environment 432: 143–158. https://doi.org/10.1016/j.scitotenv.2012.05.087

Aryal A, Harmon AC, Dugas TR (2021) Particulate matter air pollutants and cardiovascular disease: Strategies for intervention. Pharmacology & Therapeutics 223: e107890. https://doi.org/10.1016/j.pharmthera.2021.107890

Badamasi H (2017) Biomonitoring of air pollution using plants. MAYFEB Journal of Environmental Science 2: 27–39.

Basile A, Sorbo S, Conte B, Cardi M, Esposito S (2013) Ultrastructural changes and Heat Shock Proteins 70 induced by atmospheric pollution are similar to the effects observed under in vitro heavy metals stress in Conocephalum conicum (Marchantiales – Bryophyta). Environmental Pollution 182: 209–216. https://doi.org/10.1016/j.envpol.2013.07.014

Belloeil C, Jouannais P, Malfaisan C, Reyes Fernández R, Lopez S, Navarrete Gutierrez DM, Maeder-Pras S, Villanueva P, Tisserand R, Gallopin M, Alfonso-Gonzalez D, Fuentes Marrero IM, Muller S, Invernon V, Pillon Y, Echevarria G, Berazaín Iturralde R, Merlot S (2021) The X-ray fluorescence screening of multiple elements in herbarium specimens from the Neotropical region reveals new records of metal accumulation in plants. Metallomics: Integrated Biometal Science 13: mfab045. https://doi.org/10.1093/mtomcs/mfab045

Blagnytė R, Paliulis D (2010) Research into heavy metals pollution of Atmosphere applying moss as bioindicator: A literature review. Environmental Research, Engineering and Management 54: 26–33.

Bounoua L, Nigro J, Zhang P, Thome K, Lachir A (2018) Mapping urbanization in the United States from 2001 to 2011. Applied Geography 90: 123–133. https://doi.org/10.1016/j.apgeog.2017.12.002

Brook RD, Rajagopalan S, Pope CA, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC, Whitsel L, Kaufman JD (2010) Particulate Matter Air Pollution and Cardiovascular Disease: An Update to the Scientific Statement From the American Heart Association. Circulation 121: 2331–2378. https://doi.org/10.1161/CIR.0b013e3181dbece1

Chen B, Kan H (2008) Air pollution and population health: A global challenge. Environmental Health and Preventive Medicine 13: 94–101. https://doi.org/10.1007/s12199-007-0018-5

Fernández JA, Boquete MT, Carballeira A, Aboal JR (2015) A critical review of protocols for moss biomonitoring of atmospheric deposition: Sampling and sample preparation. The Science of the Total Environment 517: 132–150. https://doi.org/10.1016/j.scitotenv.2015.02.050

Fuller R, Landrigan PJ, Balakrishnan K, Bathan G, Bose-O'Reilly S, Brauer M, Caravanos J, Chiles T, Cohen A, Corra L, Cropper M, Ferraro G, Hanna J, Hanrahan D, Hu H, Hunter D, Janata G, Kupka R, Lanphear B, Lichtveld M, Martin K, Mustapha A, Sanchez-Triana E, Sandilya K, Schaefli L, Shaw J, Seddon J, Suk W, Téllez-Rojo MM, Yan C (2022) Pollution and health: A progress update. The Lancet. Planetary Health 6: e535–e547. https://doi.org/10.1016/S2542-5196(22)00090-0

Gerson U (1982) Bryophytes and invertebrates. In: Smith AJE (Ed.) Bryophyte Ecology. Springer Netherlands, Dordrecht, 291–332. https://doi.org/10.1007/978-94-009-5891-3_9

Godzik B (2020) Use of bioindication methods in national, regional and local monitoring in Poland—changes in the air pollution level over several decades. Atmosphere 11: 143. https://doi.org/10.3390/atmos11020143

Hamanaka RB, Mutlu GM (2018) Particulate matter air pollution: Effects on the cardiovascular system. Frontiers in Endocrinology 9: e680. https://doi.org/10.3389/fendo.2018.00680

Harmens H, Norris DA, Steinnes E, Kubin E, Piispanen J, Alber R, Aleksiayenak Y, Blum O, Coşkun M, Dam M, De Temmerman L, Fernández JA, Frolova M, Frontasyeva M, González-Miqueo L, Grodzińska K, Jeran Z, Korzekwa S, Krmar M, Kvietkus K, Leblond S, Liiv S, Magnússon SH, Maňkovská B, Pesch R, Rühling Å, Santamaria JM, Schröder W, Spiric Z, Suchara I, Thöni L, Urumov V, Yurukova L, Zechmeister HG (2010) Mosses as biomonitors of atmospheric heavy metal deposition: Spatial patterns and temporal trends in Europe. Environmental Pollution 158: 3144–3156. https://doi.org/10.1016/j.envpol.2010.06.039

Huttunen K-L, Mykrä H, Oksanen J, Astorga A, Paavola R, Muotka T (2017) Habitat connectivity and in-stream vegetation control temporal variability of benthic invertebrate communities. Scientific Reports 7: e1448. https://doi.org/10.1038/s41598-017-00550-9

Khomenko S, Cirach M, Pereira-Barboza E, Mueller N, Barrera-Gómez J, Rojas-Rueda D, De Hoogh K, Hoek G, Nieuwenhuijsen M (2021) Premature mortality due to air pollution in European cities: A health impact assessment. The Lancet. Planetary Health 5: e121–e134. https://doi.org/10.1016/S2542-5196(20)30272-2

Koz B, Cevik U, Ozdemir T, Duran C, Kaya S, Gundogdu A, Celik N (2008) Analysis of mosses along Sarp-Samsun highway in Turkey. Journal of Hazardous Materials 153: 646–654. https://doi.org/10.1016/j.jhazmat.2007.09.008

Landrigan PJ, Fuller R, Acosta NJR, Adeyi O, Arnold R, Basu N (Nil), Baldé AB, Bertollini R, Bose-O'Reilly S, Boufford JI, Breysse PN, Chiles T, Mahidol C, Coll-Seck AM, Cropper ML, Fobil J, Fuster V, Greenstone M, Haines A, Hanrahan D, Hunter D, Khare M, Krupnick A, Lanphear B, Lohani B, Martin K, Mathiasen KV, McTeer MA, Murray CJL, Ndahimananjara JD, Perera F, Potočnik J, Preker AS, Ramesh J, Rockström J, Salinas C, Samson LD, Sandilya K, Sly PD, Smith KR, Steiner A, Stewart RB, Suk WA, Van Schayck OCP, Yadama GN, Yumkella K, Zhong M (2018) The lancet commission on pollution and health. Lancet 391: 462–512. https://doi.org/10.1016/S0140-6736(17)32345-0

Liang D, Shi L, Zhao J, Liu P, Sarnat JA, Gao S, Schwartz J, Liu Y, Ebelt ST, Scovronick N, Chang HH (2020) Urban air pollution may enhance COVID-19 case-fatality and mortality rates in the United States. The Innovation 1: e100047. https://doi.org/10.1016/j.xinn.2020.100047

Lindo Z, Nilsson M, Gundale MJ (2013) Bryophyte-cyanobacteria associations as regulators of the northern latitude carbon balance in response to global change. Global Change Biology 19: 2022–2035. https://doi.org/10.1111/gcb.12175

Liu G, Chen T, Cui J, Zhao Y, Li Z, Liang W, Sun J, Liu Z, Xiao T (2023) Trace Metal(loid) migration from road dust to local vegetables and tree tissues and the bioaccessibility-based health risk: Impacts of vehicle operation-associated emissions. International Journal of Environmental Research and Public Health 20: e2520. https://doi.org/10.3390/ijerph20032520

Manisalidis I, Stavropoulou E, Stavropoulos A, Bezirtzoglou E (2020) Environmental and health impacts of air pollution: A review. Frontiers in Public Health 8: 1–14. https://doi.org/10.3389/fpubh.2020.00014

Mayer PM, Moran KD, Miller EL, Brander SM, Harper S, Garcia-Jaramillo M, Carrasco-Navarro V, Ho KT, Burgess RM, Thornton Hampton LM, Granek EF, McCauley M, McIntyre JK, Kolodziej EP, Hu X, Williams AJ, Beckingham BA, Jackson ME, Sanders-Smith RD, Fender CL, King GA, Bollman M, Kaushal SS, Cunningham BE, Hutton SJ, Lang J, Goss HV, Siddiqui S, Sutton R, Lin D, Mendez M (2024) Where the rubber meets the road: Emerging environmental impacts of tire wear particles and their chemical cocktails. The Science of the Total Environment 927: e171153. https://doi.org/10.1016/j.scitotenv.2024.171153

Messager ML, Davies IP, Levin PS (2021) Development and validation of in-situ and laboratory X-ray fluorescence (XRF) spectroscopy methods for moss biomonitoring of metal pollution. MethodsX 8: e101319. https://doi.org/10.1016/j.mex.2021.101319

Mohsen M, Ahmed MB, Zhou JL (2018) Particulate matter concentrations and heavy metal contamination levels in the railway transport system of Sydney, Australia. Transportation Research Part D, Transport and Environment 62: 112–124. https://doi.org/10.1016/j.trd.2018.02.015

Newby DE, Mannucci PM, Tell GS, Baccarelli AA, Brook RD, Donaldson K, Forastiere F, Franchini M, Franco OH, Graham I, Hoek G, Hoffmann B, Hoylaerts MF, Künzli N, Mills N, Pekkanen J, Peters A, Piepoli MF, Rajagopalan S, Storey RF (2015) Expert position paper on air pollution and cardiovascular disease. European Heart Journal 36: 83–93. https://doi.org/10.1093/eurheartj/ehu458

Nieuwenhuijsen MJ (2020) Urban and transport planning pathways to carbon neutral, liveable and healthy cities; A review of the current evidence. Environment International 140: e105661. https://doi.org/10.1016/j.envint.2020.105661

Pearson J, Wells DM, Seller KJ, Bennett A, Soares A, Woodall J, Ingrouille MJ (2000) Traffic exposure increases natural$^{15}$ N and heavy metal concentrations in mosses. The New Phytologist 147: 317–326. https://doi.org/10.1046/j.1469-8137.2000.00702.x

Postiglione A, Dentato M, Maresca V, Basile A (2025) Liverworts as heavy metals and metalloids bioindicators: Bioaccumulation, biological responses and future applications in environmental biomonitoring. Plant Biosystems •••: 1–13. https://doi.org/10.1080/11263504.2025.2525872

Rieley JO, Richards PW, Bebbington ADL (1979) The ecological role of bryophytes in a north wales woodland. Journal of Ecology 67: e497. https://doi.org/10.2307/2259109

Roblin B, Aherne J (2020) Moss as a biomonitor for the atmospheric deposition of anthropogenic microfibres. The Science of the Total Environment 717: e136973. https://doi.org/10.1016/j.scitotenv.2020.136973 [DOI] [PubMed] [Google Scholar]

Ruehling A, Tyler G (1968) Ecological approach to the lead problem. ETDEWEB 121: 1–3. https://www.osti.gov/etdeweb/biblio/5556070 [February 7, 2025]

Ruklani S, Rubasinghe SCK, Jayasuriya G (2021) A review of frameworks for using bryophytes as indicators of climate change with special emphasis on Sri Lankan bryoflora. Environmental Science and Pollution Research International 28: 60425–60437. https://doi.org/10.1007/s11356-021-16588-2

Sharma SB, Jain S, Khirwadkar P, Kulkarni S (2013) The effects of air pollution on the environment and human health. Indian Journal of Research in Pharmacy and Biotechnology 1(3): 391–396.

Shleag AM, Shiha FA, Aborba NA (2024) Environmental and air pollution's impact on health: Challenges and opportunities. Zenodo. https://doi.org/10.5281/ZENODO.13363655

Šoltés R, Gregušková E (2013) Accumulation characteristics of some elements in the *Moss polytrichum commune* (Bryophytes) based on XRF spectrometry. Journal of Environmental Protection 04: 522–528. https://doi.org/10.4236/jep.2013.46061

Thomas M (1961) Effects of air pollution on plants. World Health Organization Monograph Series 46: 233–278.

Travaglio M, Yu Y, Popovic R, Selley L, Leal NS, Martins LM (2021) Links between air pollution and COVID-19 in England. Environmental Pollution 268: e115859. https://doi.org/10.1016/j.envpol.2020.115859

Urošević MA, Lazo P, Stafilov T, Nečemer M, Andonovska KB, Balabanova B, Hristozova G, Papagiannis S, Stihi C, Suljkanović M, Špirić Z, Vassilatou V, Vogel-Mikuš K (2023) Active biomonitoring of potentially toxic elements in urban air by two distinct moss species and two analytical techniques: A pan-Southeastern European study. Air Quality, Atmosphere & Health 16: 595–612. https://doi.org/10.1007/s11869-022-01291-z

Vegetation ICP (2020) Monitoring of atmospheric deposition of heavy metals, nitrogen and POPs in Europe using Bryophytes. United Nations Economic Commission for Europe Convention on Long-Range Transboundary Air Pollution. https://icpvegetation.ceh.ac.uk/sites/default/files/ICP%20Vegetation%20moss%20monitoring%20manual%202020.pdf [August 1, 2025]

Weinberger M, Schillaci L, Rumancev C, Bretschneider T, Rosenhahn A, Lang I, Probst A (2025) Synchrotron source X-ray experiments reveal element-specific distributions of heavy metals in moss leaves. Program & Book of Abstracts. BOKU University, Vienna, Austria, 100. https://boku.ac.at/fileadmin/data/H03000/H83000/H83100/files/ATSPB/ATSPB_2025_BoA.pdf [August 4, 2025]